# Oncology Clinical Pathways Esophageal Cancer

September 2022 - V1.2022








#### **Table of Contents**

Stage 1A T1aN0M0 Adenocarcinoma or Squamous Cell Carcinoma	3
Stage I T1bN0M0 Adenocarcinoma or Squamous Cell Carcinoma	4
Stage T1N1(IIA), T2N0(IIB), T2N1 (III), T3 N0-1(III), T4aN0-1(III), T1-4AN2(IVA), T1-3 N3 (IVA) Adenocarcinoma	5
Stage T2-T3 N0 (IB-IIB), T1-T2 N1(IIB), T1-2 N2 (IIIA), T3N1 (IIIA), T4aN0-1 (IIIA), T3N2(IIIB), T4aN2 (IVA) Non-Cervical Squamous Cell Carcinoma	6
Stage IVB Adenocarcinoma Molecular Testing.	7
Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Positive First Line.	8
Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Positive Second Line.	9
Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Negative First Line.	10
Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Negative Second Line.	11
Stage IVB Adenocarcinoma Any T, Any N, M1 MSI High	12
Stage IVB Squamous Cell Carcinoma Any T, Any N, M1 First Line.	
Stage IVB Squamous Cell Carcinoma Any T, Any N, M1 Second Line	14
Cervical Squamous or Unresectable Localized.	15





#### <u>Esophageal Cancer – Stage 1A T1aN0M0</u> <u>Adenocarcinoma or Squamous Cell Carcinoma</u>




- <sup>a</sup> **Staging** confirmed during Endoscopic Mucosal Resection (EMR) or Endoscopic Submucosal Dissection (ESD)
- <sup>b</sup> Multidisciplinary discussion includes surgery, radiation oncology, and medical oncology







#### <u>Esophageal Cancer – Stage I T1bN0M0</u> <u>Adenocarcinoma or Squamous Cell Carcinoma</u>



Clinical trial(s) always considered on pathway.

- <sup>a</sup> Multidisciplinary discussion confirm stage
- <sup>b</sup> High risk features include lymphovascular invasion, poorly differentiated tumors, tumor size ≥ 2cm, SM1, SM2, or positive margin on EMR


**EMR** Endoscopic Mucosal Resection







## Esophageal Cancer – Stage T1N1(IIA), T2N0(IIB), T2N1 (III), T3 N0-1(III), T4aN0-1(III), T1-4AN2(IVA), T1-3 N3 (IVA) Adenocarcinoma



Clinical trial(s) always considered on pathway

<sup>a</sup> Multidisciplinary discussion includes surgery, radiation oncology, and medical oncology

Surgical candidate discussion with or evaluation by a surgeon

Carboplatin and paclitaxel with radiation preferred treatment for predominantly esophageal tumors

Perioperative chemotherapy preferred treatment for predominantly gastric tumors or patient is not a candidate for radiation

e FLOT4 candidates include fit patients with ECOG 0-1 due to expected Grade 3/4 toxicities of neutropenia, infection, diarrhea and peripheral neuropathy


f Qualify for immune checkpoint inhibitor patient without active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent > 10mg/day) or prior allogeneic HSCT/solid organ transplant







#### Esophageal Cancer – Stage T2-T3 N0 (IB-IIB), T1-T2 N1(IIB), T1-2 N2 (IIIA), T3N1 (IIIA), T4aN0-1 (IIIA), T3N2(IIIB), T4aN2 (IVA) **Non-Cervical Squamous Cell Carcinoma**



Clinical trial(s) always considered on pathway

<sup>a</sup> Multidisciplinary discussion includes surgery, radiation oncology, and medical oncology

Surgical candidate discussion with or evaluation by a surgeon


Qualify for immune checkpoint inhibitor patient without active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent >10mg/day) or prior allogeneic HSCT/solid organ transplant







#### <u>Esophageal Cancer – Stage IVB Adenocarcinoma Molecular Testing</u>



Clinical trial(s) always considered on pathway.

<sup>a</sup> MSI or MMR NGS or IHC


<sup>b</sup> **HER2 IHC** if results 2+, HER2 IHC with reflex to FISH required (in-situ hybridization)







#### <u>Esophageal Cancer – Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Positive First Line</u>




- <sup>a</sup> HER2 Positive considered IHC score +3 or an IHC score of +2 and FISH/ISH positive
- Assess nutritional status and consider palliative stent or other nutritional support modalities when clinically appropriate
- Candidate for cytotoxic chemotherapy consider if patient can tolerate a platinum- and fluoropyrimidine-based doublet
- d Qualify for trastuzumab or biosimilar patient with HER2-positive disease and no clinically significant CV disease (defined as LVEF< 50%, MI within prior 6 months, symptomatic CHF (NYHA class II to IV), unstable angina or cardiac arrhythmia requiring therapy)
- Radiation Oncology consider palliative radiation when clinically appropriate
- Qualify for immune checkpoint inhibitor patient without active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent >10mg/day) or prior allogeneic HSCT/solid organ transplant
- Pembrolizumab for two years







### <u>Esophageal Cancer – Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Positive Second Line</u>










<sup>&</sup>lt;sup>a</sup> Candidate for fam-trastuzumab deruxtecan received trastuzumab in the first-line setting; baseline LVEF ≥ 50% and/or no clinically significant cardiac disease (defined as LVEF < 50%, MI within prior 6 months, symptomatic CHF (NYHA class II to IV), unstable angina or cardiac arrhythmia requiring therapy); no ILD or pneumonitis; ANC ≥ 1500/mm³

b Candidate for ramucirumab received fluoropyrimidine and platinum agent in the first-line setting; ECOG PS 0-2; ANC ≥ 1500/mm³. Note: Due to anti-VEGF effects patients with the following should <u>not</u> receive ramucirumab: non-healing wound/fracture, major surgery in prior 4 weeks, bleeding disorder or coagulopathy, recent history of GI perforation, unstable cardiac condition (uncontrolled HTN, arterial thromboembolism, symptomatic CHF (NYHA II-IV) or arrhythmia), or active cocaine use

### <u>Esophageal Cancer – Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Negative First Line</u>




- Assess nutritional status consider palliative stent or other nutritional support modalities when clinically appropriate
- <sup>b</sup> Assess Palliative Care consider palliative radiation when clinically appropriate
- <sup>c</sup> Candidate for cytotoxic chemotherapy consider if patient can tolerate a platinum- and fluoropyrimidine-based doublet
- <sup>d</sup> Qualify for immune checkpoint inhibitor no active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent >10mg/day) or prior allogeneic HSCT/solid organ transplant







### <u>Esophageal Cancer – Stage IVB Adenocarcinoma Any T, Any N, M1 MSS HER2 Negative Second Line</u>



Clinical trial(s) always considered on pathway.


<sup>a</sup> Candidate for ramucirumab received fluoropyrimidine and platinum agent in the first-line setting; ECOG PS 0-2; ANC ≥ 1500/mm<sup>3</sup>; due to anti-VEGF effects patients with the following should <u>not</u> receive ramucirumab: non-healing wound/fracture, major surgery in prior 4 weeks, bleeding disorder or coagulopathy, recent history of GI perforation, unstable cardiac condition (uncontrolled HTN, arterial thromboembolism, symptomatic CHF (NYHA II-IV) or arrhythmia), or active cocaine use







### Esophageal Cancer – Stage IVB Adenocarcinoma Any T, Any N, M1 MSI High










<sup>&</sup>lt;sup>a</sup> Assess nutritional status consider palliative stent or other nutritional support modalities when clinically appropriate

<sup>&</sup>lt;sup>b</sup> Qualify for immune checkpoint inhibitor patient without active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent >10mg/day) or prior allogeneic HSCT/solid organ transplant

## <u>Esophageal Cancer – Stage IVB Squamous Cell Carcinoma</u> <u>Any T, Any N, M1 First Line</u>



d Radiation Oncology consider palliative or metastasis-directed radiation when clinically appropriate








<sup>&</sup>lt;sup>a</sup> Assess nutritional status consider palliative stent or other nutritional support modalities when clinically appropriate

<sup>&</sup>lt;sup>b</sup> Qualify for immune checkpoint inhibitor patient without active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent >10mg/day) or prior allogeneic HSCT/solid organ transplant

<sup>&</sup>lt;sup>c</sup> Candidate for cytotoxic chemotherapy consider if patient can tolerate a platinum- and fluoropyrimidine-based doublet

### <u>Esophageal Cancer – Stage IVB Squamous Cell Carcinoma</u> <a href="mailto:Any T, Any N, M1 Second Line">Any T, Any N, M1 Second Line</a>



<sup>a</sup> Qualify for immune checkpoint inhibitor patient without active autoimmune disease, primary immune deficiency, concurrent immunosuppression (including prednisone equivalent >10mg/day) or prior allogeneic HSCT/solid organ transplant


<sup>b</sup> Pembrolizumab chosen due to FDA approval and patient convenience







#### <u>Esophageal Cancer – Cervical Squamous or Unresectable Localized</u>



Clinical trial(s) always considered on pathway.

<sup>a</sup> Multidisciplinary discussion includes surgery, radiation oncology, and medical oncology; for mid-esophageal tumors, consider referral to Pulmonary to assess tracheal invasion







#### **Questions?**

Contact VHAOncologyPathways@va.gov





